

Key Beef Finisher 12 % R

9007503

MEDICATED

Cattle fed in confinement for slaughter: For improved feed efficiency.

ACTIVE DRUG INGREDIENT
Monensin Sodium 33 g/ton

A Complete Feed for Beef Cattle

GUARANTEED ANALYSIS

Crude Protein		(Min)		12.30 %
Crude Fat		(Min)		3.00 %
Crude Fiber		(Max)		5.50 %
Calcium (Ca)	(Min)	0.51	(Max)	1.21 %
Phosphorus (P)		(Min)		0.29 %

INGREDIENTS

Grain Products, Plant Protein Products, Molasses Products, Roughage Products, Processed Grain By-Products, Calcium Carbonate, Salt, Magnesium Oxide, Propionic acid, Zinc Sulfate, Manganous Oxide, Ferrous Sulfate, Copper Sulfate, Sodium Selenite, Vitamin E Supplement, Vitamin A Acetate, Vitamin D3 Supplement, Ethylenediamine Dihydroiodide, Cobalt Carbonate.

FEEDING DIRECTIONS

Feed Key Beef Finisher 12 % R with good quality forages and/or roughages to meet the nutritional requirements of Beef Cattle. Free choice salt may be provided in a mineral feeder. For a more specific feeding program, please contact your Keystone Mills Feed Consultant.

CAUTION: Do not feed supplemental selenium in excess of 0.3 ppm in the total nutritional diet.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: Feed continuously in complete feed at a rate of 50 to 480 milligrams of monensin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton.



Romulus, NY 14541 315-549-8226

NET WT. 50 lbs (22.68 kg)

Lot#:01-0000296190-188

RED

rev: 02/09/17