

Key Goat Grower 16% Pellet R

9303243

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christensenii*, and *E. ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin Sodium 25 g/ton

Complete Feed for Goats

GUARANTEED ANALYSIS

Crude Protein (Min) 16.20 %
* (This Includes not more than 0.8 % equivalent non-protein nitrogen.)

Crude Fat		(Min)	3.80 %
Crude Fiber		(Max)	10.00 %
Calcium (Ca)	(Min)	0.82	(Max) 1.22 %
Phosphorus (P)		(Min)	0.50 %
Salt (NaCl)	(Min)	0.32	(Max) 0.72 %

INGREDIENTS

Processed Grain By-Products, Grain Products, Roughage Products, Plant Protein Products, Forage Products, Molasses Products, Calcium Carbonate, Salt, Ammonium Chloride, Maxi Bond, Potassium Magnesium Sulfate, Dicalcium Phosphate, Propionic acid, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Selenium Yeast, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Vitamin E, Choline Chloride, Copper Sulfate, Niacin, Copper Amino Acid Complex, Sodium Selenite, Thiamine Mononitrate, Ethylenediamine Dihydroiodide, Vitamin A, Vitamin D3, d-Calcium Pantothenate, Biotin, Folic Acid, Vitamin B12, Cobalt Carbonate, Riboflavin, Pyridoxine Hydrochloride, Mineral Oil, Yeast Culture, Yeast Extract, Hydrolyzed Yeast.

FEEDING DIRECTIONS

Feed Key Goat Grower 16% Pellet R with good quality forages and/or roughages to meet the nutritional requirements of growing goats. Free choice salt may be provided. Supply plenty of fresh clean water at all times. For a more specific feeding program, please contact your Keystone Mills Feed Consultant.

CAUTION: Do not feed supplemental selenium in excess of 0.3 ppm in the total nutritional diet.

CAUTION: USE AS DIRECTED

THIS FEED SHOULD BE USED ONLY IN ACCORDANCE WITH THE RECOMMENDATIONS ON THIS FEED TAG

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is safe for use in goats only. Consumption by unapproved species may result in toxic reactions.

LIMITATIONS: Feed continuously. Do not feed to lactating goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the level of Monensin recommended in the feeding direction, as reduced average daily gains may result.

WARNING: Must Observe feeding directions on label.



Romulus, NY 14541 315-549-8226

NET WT. 50 lbs (22.68 kg)

Lot#: 01-0000296487-193

rev: 07/23/19