

Key Kid Starter 18% Pellet R

9303253

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christenseni*, and *E. ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin Sodium 30 g/ton

Complete Feed for Goats

GUARANTEED ANALYSIS

Crude Protein (Min) 18.20 %
* (This Includes not more than 0.8 % equivalent non-protein nitrogen.)

Crude Fat	(Min)	3.50 %
Crude Fiber	(Max)	10.00 %
Calcium (Ca)	(Min) 0.80	(Max) 1.50 %
Phosphorus (P)	(Min)	0.49 %
Salt (NaCl)	(Min) 0.28	(Max) 0.98 %

INGREDIENTS

Processed Grain By-Products, Grain Products, Plant Protein Products, Roughage Products, Forage Products, Calcium Carbonate, Molasses Products, Salt, Ammonium Chloride, Maxi Bond, Potassium Magnesium Sulfate, Propionic acid, Magnesium Oxide, Dicalcium Phosphate, Sodium Selenite, Mineral Oil, Manganous Hydroxy chloride, Zinc Hydrochloride, Vitamin E Supplement, Basic Copper Chloride, Vitamin A Acetate, Vit D3 Supplement, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Natural Flavors., Zinc Sulfate, Manganous Oxide, Ferrous Sulfate, Vitamin D3 Supplement, Vitamin B12 Supplement, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Yeast Culture, Yeast Extract, Hydrolyzed Yeast.

FEEDING DIRECTIONS Feed Key Kid Starter 18% Pellet R with good quality forages and/or roughages to meet the nutritional requirements of growing goats. Free choice salt may be provided. Supply plenty of fresh clean water at all times. For a more specific feeding program, please contact your Keystone Mills Feed Consultant. **CAUTION:** Do not feed supplemental selenium in excess of 0.3 ppm in the total nutritional diet. **CAUTION: USE AS DIRECTED THIS FEED SHOULD BE USED ONLY IN ACCORDANCE WITH THE RECOMMENDATIONS ON THIS FEED TAG** **CAUTION:** Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is safe for use in goats only. Consumption by unapproved species may result in toxic reactions. **LIMITATIONS:** Feed continuously. Do not feed to lactating goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the level of Monensin recommended in the feeding direction, as reduced average daily gains may result. **WARNING:** Must Observe feeding directions on label.

Manufactured By
KEYSTONE MILLS
Serving Agriculture Since 1894
Romulus, NY 14541 315-549-8226

NET WT. 50 lbs (22.68 kg)

Lot#:0000295394-176

rev: 10/02/18

ORANGE