

Key Calf Starter

18% R FC

9512053

MEDICATED

For the prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii* in calves (excluding veal calves).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)..... 35 g/ton

Feed for Calves from 3 Days to 3 Months of Age

GUARANTEED ANALYSIS

Crude Protein		(Min)	17.80 %
Crude Fat		(Min)	4.00 %
Crude Fiber		(Max)	5.00 %
Calcium (Ca)	(Min)	0.50	(Max) 1.20 %
Phosphorus (P)		(Min)	0.50 %
Salt (NaCl)	(Min)	0.28	(Max) 0.98 %

INGREDIENTS: Grain Products, Plant Protein Products, Processed Grain By-Products, Molasses Products, Calcium Carbonate, Vegetable Oil, Dicalcium Phosphate, Salt, Animal Fat, Propionic acid, Maxi Bond, Roughage Products, Natural and Artificial Flavors, Vitamin E Supplement, Ferrous Sulfate, Hydrated Sodium Silico Aluminate, Vitamin D3 Supplement, Zinc Sulfate, Vegetable Fat, Silicon Dioxide, Biotin, Zinc Proteinate, Manganese Proteinate, Manganese Oxide, Copper Sulfate, Sodium Selenite, Copper Proteinate, Vitamin A Acetate, Niacin Supplement, Ethylenediamine Dihydroiodide, d-Calcium Pantothenate, Vitamin B12 Supplement, Folic Acid, Riboflavin Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, Cobalt Carbonate, Menadione Nicotinamide Bisulfite.

FEEDING DIRECTIONS: Feed to provide 0.14 mg to 0.42 mg of monensin per pound of body weight. A 150 lb calf must consume 1.4 lb to 4.2 lbs of Calf Starter daily. Provide 2 to 3 gallons of fresh clean water daily in addition to milk or milk replacer. For further information, please contact your Keystone Mills Dairy Consultant.

CAUTION: Do not feed supplemental selenium in excess of 0.3 ppm in the total nutritional diet. **CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **WARNING:** A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

Manufactured By
KEYSTONE MILLS
Serving Agriculture Since 1894
Romulus, NY 14541 315-549-8226

NET WT. 50 lbs (22.68 kg)

Lot#:01-0000256789-186

rev: 08/29/17